

SQUARES OF CHARACTERS AND FINITE GROUPS

EDITH ADAN-BANTE

ABSTRACT. Let G be a group of odd order and χ be a complex irreducible character. Then there exists a unique character $\chi^{(2)} \in \text{Irr}(G)$ such that $[\chi^2, \chi^{(2)}]$ is odd. Also, there exists a unique character $\psi \in \text{Irr}(G)$ such that $[\psi^2, \chi]$ is odd.

1. INTRODUCTION

Let G be a finite group. Denote by $\text{Irr}(G)$ the set of irreducible complex characters of G . Through this work, we use the notation of [3]. In addition, we are going to denote by $\text{Lin}(G) = \{\lambda \in \text{Irr}(G) \mid \lambda(1) = 1\}$ the set of linear characters.

Let ϑ be a generalized character of G , i.e $\vartheta \in \mathbb{Z}[\text{Irr}(G)]$. We define $\vartheta^{(2)}$ by $\vartheta^{(2)}(g) = \vartheta(g^2)$ for all $g \in G$. By Theorem 4.5 of [3], we have that $\vartheta^{(2)}$ is also a generalized character. Also we can check that if $\vartheta \in \text{Irr}(G)$ and G is of odd order, then $\vartheta^{(2)} \in \text{Irr}(G)$ (see Exercise 4.5 of [3]).

Let G be a group of odd order and $\lambda \in \text{Lin}(G)$. Since $\lambda^2 \in \text{Lin}(G)$, λ^2 has a unique irreducible constituent that appears with odd multiplicity, namely λ^2 , and such character has the same degree as λ . Since $|G|$ is odd and $\lambda \in \text{Lin}(G)$, we can check that there exists $\mu \in \text{Lin}(G)$ such that λ is the unique irreducible constituent that appears with odd multiplicity in the character μ^2 . So we can find an irreducible character that is “the square” (“the square root”) given any linear character. For any character $\chi \in \text{Irr}(G)$, where χ is not necessarily linear, the following result implies that $\chi^{(2)} \in \text{Irr}(G)$ is the unique irreducible character of G that appears with odd multiplicity in the character χ^2 . So we can regard $\chi^{(2)}$ as “the square” of χ .

Theorem A. *Let G be a finite group of odd order. The map $\vartheta \mapsto \vartheta^{(2)}$ is an automorphism of the ring $\mathbb{Z}[\text{Irr}(G)]$. Also, if $\vartheta \in \mathbb{Z}[\text{Irr}(G)]$ then*

$$(1.1) \quad \vartheta^2 = \vartheta^{(2)} + 2\Gamma$$

for some $\Gamma \in \mathbb{Z}[\text{Irr}(G)]$.

We now study squares of irreducible characters of prime degree.

Theorem B. *Let G be a finite group and $\chi \in \text{Irr}(G)$ be a monomial character such that $\chi(1) = 2$. Then one of the following holds*

- (i) $\chi^2 = \lambda_1 + \lambda_2 + \theta$, for some $\lambda_1, \lambda_2 \in \text{Lin}(G)$ and some $\theta \in \text{Irr}(G)$ with $\theta(1) = 2$.
- (ii) $\chi^2 = \sum_{i=1}^4 \lambda_i$ for some distinct linear characters $\lambda_1, \lambda_2, \lambda_3$ and λ_4 .

Date: 2004.

1991 Mathematics Subject Classification. 20c15.

Key words and phrases. Products of characters, groups of odd order, irreducible constituents.

Let Θ be a character of G . Denote by $\eta(\Theta)$ the number of distinct irreducible constituents of the character Θ . In Theorem 2.3 is proved that if G is a supersolvable group and $\chi \in \text{Irr}(G)$ with $\chi(1) = 2^n$ for some integer $n > 0$, then $\eta(\chi^2) \geq 2$, that is χ^2 is the sum of at least two distinct irreducible characters.

Theorem C. *Let p be an odd prime number. Let G be a finite nilpotent group and $\chi \in \text{Irr}(G)$ with $\chi(1) = p$. Then all the irreducible constituents of χ^2 have degree p and one of the following holds:*

- (i) $\eta(\chi^2) = 1$. Also $\chi^2 = p\chi^{(2)}$ and χ vanishes outside $\mathbf{Z}(\chi)$.
- (ii) $\eta(\chi^2) = \frac{p+1}{2}$. Furthermore

$$\chi^2 = \chi^{(2)} + 2 \sum_{i=1}^{\frac{p-1}{2}} \theta_i$$

where $\theta_1, \theta_2, \dots, \theta_{\frac{p-1}{2}}$ are distinct irreducible characters of G .

Let G be a nilpotent group and $\chi \in \text{Irr}(G)$ with $\chi(1) = p$. By Theorem B and Theorem C we have that if p is a prime number, then there are exactly 2 possible values for $\eta(\chi^2)$. Assume now that $\chi(1) = p^n$ for a fixed prime p . The author wonders for what integer $n > 1$, the number of possible values for $\eta(\chi^2)$ really does depend upon the prime p .

In Remark 2.10, examples are provided, showing that Theorem C may not remain true if G is only a supersolvable group of odd order.

Acknowledgment. I thank Professor Everett C. Dade for very useful conversations and emails. I also thank Professor Avinoam Mann and the referee for their corrections and suggestions to improve this note.

2. PROOFS

Proof of Theorem A. Exercise 4.7 of [3] implies (1.1). It remains then to prove that the map $\vartheta \mapsto \vartheta^{(2)}$ is a bijection.

Since G has odd order, given any character f and any element $g \in G$, there exists a unique element $f'(g)$ over the complex such that $(f'(g))^2 = f(g)$. Let f' be the function defined as $(f'(g))^2 = f(g)$ for all $g \in G$. If f is any character, so is f' and $(f')^2 = f$. That shows that the map $f \mapsto f'$ is onto on the set of characters, and therefore one to one. We can then extend this to generalized characters. $\square$

Let G be the dihedral group of order 8 and $\chi \in \text{Irr}(G)$ with $\chi(1) = 2$. We can check that

$$\chi^2 = \sum_{\lambda \in \text{Lin}(G)} \lambda.$$

Thus for a group G of even order and $\chi \in \text{Irr}(G)$, there may not exist a unique character that appears with odd multiplicity in χ^2 .

Proof of Theorem B. Without loss of generality, we may assume that χ is a faithful character. Since χ is monomial, there exists a subgroup N of G and a linear character λ of N such that $\lambda^G = \chi$. Since $\chi(1) = 2$, then $|G : N| = 2$ and thus N is a normal subgroup of G . Since χ is faithful and $\chi(1) = 2$, we have that N is an abelian subgroup of G . Fix $g \in G \setminus N$. By Clifford theory we have that $\lambda^g \neq \lambda$ and $\chi_N = \lambda + \lambda^g$. Thus

$$(\chi^2)_N = \lambda^2 + (\lambda^g)^2 + 2\lambda\lambda^g.$$

Observe that $\lambda\lambda^g$ is a G -invariant character of N . Since $|G : N| = 2$, it follows that $\lambda\lambda^g$ extends to G .

Let $\mu \in \text{Lin}(G)$ be an extension of $\lambda\lambda^g$. Observe that $\chi\nu = \chi$ for any $\nu \in \text{Lin}(G/N)$. Thus χ^2 has at least 2 linear constituents, namely the two distinct extensions of the character $\lambda\lambda^g$. By Exercise 4.12 of [3], we have that $[\chi^2, \theta] \leq \theta(1)$ for any $\theta \in \text{Irr}(G)$. Thus χ^2 has at least 3 distinct irreducible constituents, i.e. $\eta(\chi^2) \geq 3$. Since $\chi^2(1) = 4$, either $\eta(\chi^2) = 3$ or $\eta(\chi^2) = 4$.

If $\eta(\chi^2) = 3$ then χ^2 has 2 linear constituents and one irreducible constituent of degree 2 and (i) holds.

If $\eta(\chi^2) = 4$ then all the irreducible constituents of χ^2 are linear and (ii) holds. $\square$

Remark 2.1. Let $\text{SL}(2, 3)$ be the special linear group of degree 2 over the field of 3 elements, and $\chi \in \text{Irr}(\text{SL}(2, 3))$ with $\chi(1) = 2$. We can check that $\chi^2 = \lambda + \theta$, where $\theta \in \text{Irr}(G)$ and $\theta(1) = 3$ and $\lambda \in \text{Lin}(\text{SL}(2, 3))$. Thus Theorem B does not remain true without the hypothesis that χ is a monomial character.

Remark 2.2. Let C_{2^n} and C_2 be cyclic subgroups of order 2^n , where $n > 1$, and 2 respectively. Set $N = C_{2^n} \times C_{2^n}$. Observe that C_2 acts on N by permutation. Let G be the semidirect product of N and C_2 . Thus $|G| = 2^{2n+1}$.

Let $(\lambda, 1_{C_{2^n}}) \in \text{Irr}(N)$ be a character such that $\lambda^2 \neq 1_{C_{2^n}}$. Set $\chi = (\lambda, 1_{C_{2^n}})^G$ and $\theta = (\lambda^2, 1_{C_{2^n}})^G$. Observe that χ and θ are irreducible characters of degree 2. We can check that

$$\chi^2 = \lambda_1 + \lambda_2 + \theta$$

where λ_1 and λ_2 are the 2 distinct extensions of the G -invariant character $(\lambda, \lambda) \in \text{Lin}(N)$.

Let G be the dihedral subgroup of order 8. We can check that χ^2 is the sum of 4 linear characters.

We conclude that for each of the cases in Theorem B, there exists a group G and a character $\chi \in \text{Irr}(G)$ satisfying the conditions in that case.

Theorem 2.3. *Let G be a finite supersolvable group of G and $\chi \in \text{Irr}(G)$. Assume that $\chi(1) = 2^n$ for some integer $n > 0$. Then $\eta(\chi^2) > 1$.*

Proof. Assume that the statement is false. Let G be a supersolvable group and $\chi \in \text{Irr}(G)$ be a character of G such that they are a minimal counterexample with respect to the order of G . So $\chi^2 = m\theta$ for some integer m , some $\theta \in \text{Irr}(G)$ and $\chi(1) = 2^n$ for some integer $n > 0$.

Let $Z = \mathbf{Z}(\chi)$ be the center of the character χ . Since $\chi^2 = m\theta$, we can check that $Z = \mathbf{Z}(\theta)$. Let $\lambda \in \text{Irr}(Z)$ be the unique linear character of Z such that $\chi_Z = \chi(1)\lambda$. Observe that $\theta_Z = \theta(1)\lambda^2$ since $\chi^2 = m\theta$. Let Y/Z be a chief factor of G and $\mu \in \text{Irr}(Y)$ be a character such that $[\chi_Y, \mu] \neq 0$. Because G is supersolvable and Y/Z is a chief factor of G , it follows that Y/Z is cyclic and therefore λ extends to Y . Thus μ is a linear character.

Let $H = G_\mu$ be the stabilizer of μ in G and $\chi_\mu \in \text{Irr}(H)$ be the Clifford correspondent of χ and μ , i.e. $\chi_\mu^G = \chi$. Since μ is a linear character, $[\chi_Y, \mu] \neq 0$ and $Y > Z = \mathbf{Z}(\chi)$, we have that H is a proper subgroup of G . Observe that $[\theta_Y, \mu^2] \neq 0$. We can check that the stabilizer G_{μ^2} of μ^2 in G contains $G_\mu = H$ and $G_{\mu^2} < G$. Let K be a subgroup of G such that $G_{\mu^2} \leq K < G$ and χ_μ^K is not a linear character. Observe such subgroup K exists since $\chi(1) > 2$ by Theorem B.

By Clifford theory we have that $\chi_\mu^K \in \text{Irr}(G)$. Since χ_μ^K lies above μ , all the irreducible constituents of $\chi_\mu^K \chi_\mu^K$ lie above μ^2 . Since $G_{\mu^2} \leq K$, by Clifford theory we can check that $\eta(\chi_\mu^K \chi_\mu^K) = \eta((\chi_\mu^K \chi_\mu^K)^G)$. Since $|K| < |G|$ and $\chi_\mu(1)$ is not a linear character, we have that $\eta(\chi_\mu^K \chi_\mu^K) > 1$. Since $(\chi_\mu^K)^G = \chi$, we can check that $\eta((\chi_\mu^K \chi_\mu^K)^G) \leq \eta(\chi\chi)$. But then $1 < \eta(\chi_\mu^K \chi_\mu^K) \leq \eta(\chi\chi) = 1$. Therefore such group G and such character χ can not exist. $\square$

Recall that the irreducible characters of supersolvable groups are monomial. The author wonders if Theorem 2.3 remains true with the weaker hypothesis that χ is monomial.

A corollary of Theorem 2.3 is the following

Corollary 2.4. *Let G be a nilpotent group and $\chi \in \text{Irr}(G)$. If χ^2 is a multiple of an irreducible, then $\chi(1)$ is an odd integer.*

Proposition 2.5. *Let G be a group of odd order and $\chi \in \text{Irr}(G)$. If $\chi(1) > 1$ then $[\chi^2, \lambda] = 0$ for all $\lambda \in \text{Lin}(G)$.*

Proof. Assume that $[\chi^2, \lambda] \neq 0$ for some $\lambda \in \text{Lin}(G)$. Since G has odd order, there exists some character $\beta \in \text{Lin}(G)$ such that $\beta^2 = \lambda$. Thus $[\chi^2, \lambda] = [\chi^2, \beta^2] = [\chi\bar{\beta}, \bar{\chi}\beta]$. Since $\chi\bar{\beta}, \bar{\chi}\beta \in \text{Irr}(G)$, it follows that

$$\chi\bar{\beta} = \bar{\chi}\beta.$$

Observe that $\bar{\chi}\beta = \overline{\chi\bar{\beta}}$. Since $\chi\bar{\beta} \in \text{Irr}(G)$ is a real character and G has odd order, it follows that $\chi\bar{\beta} = 1_G$. Thus $\chi = \beta$ and $\chi(1) = 1$. $\square$

Proposition 2.6. *Let G be a finite group and $\chi \in \text{Irr}(G)$. Assume that $\text{Ker}(\chi) \neq \mathbf{Z}(\chi)$. Then $[\chi^2, \chi] = 0$.*

Proof. Observe that $[\chi^2, \chi] = [\chi, \chi\bar{\chi}]$. Observe also that $\text{Ker}(\chi\bar{\chi}) = \mathbf{Z}(\chi)$. Thus if $[\chi, \chi\bar{\chi}] \neq 0$, then $\text{Ker}(\chi) \geq \mathbf{Z}(\chi)$. Therefore $\text{Ker}(\chi) = \mathbf{Z}(\chi)$. $\square$

Corollary 2.7. *Let G be a finite p -group, where p is a prime number, and $\chi \in \text{Irr}(G)$. If $\chi \neq 1_G$ then $[\chi^2, \chi] = 0$.*

Remark 2.8. Let D be the dihedral group of order 8 and $\chi \in \text{Irr}(D)$ be such that $\chi(1) = 2$. We can check that χ^2 is the sum of 4 linear characters. Thus Proposition 2.5 may not hold true when G has even order.

Let G be the semi-direct product $\langle \tau \rangle \rtimes \langle \sigma \rangle$, where σ and τ have orders 7 and 3, respectively, and $\sigma^\tau = \sigma^2$. We can check that any faithful irreducible character χ of G has degree 3, $\mathbf{Z}(\chi) = 1$ and $[\chi^2, \chi] = 1$. Thus Proposition 2.6 may not remain true if $\text{Ker}(\chi) = \mathbf{Z}(\chi)$.

Lemma 2.9. *Let G be a finite p -group, where p is an odd prime, and $\chi \in \text{Irr}(G)$ with $\chi(1) = p$. Then all the irreducible constituents of χ^2 have degree p and one of the following holds:*

- (i) $\eta(\chi^2) = 1$. Also $\chi^2 = p\chi^{(2)}$ and χ vanishes outside $\mathbf{Z}(\chi)$.
- (ii) $\eta(\chi^2) = \frac{p+1}{2}$. Furthermore

$$\chi^2 = \chi^{(2)} + 2 \sum_{i=1}^{\frac{p-1}{2}} \theta_i$$

where $\theta_1, \theta_2, \dots, \theta_{\frac{p-1}{2}}$ are distinct irreducible characters of G .

Proof. By working with the group $G/\text{Ker}(\chi)$, without loss of generality we may assume that $\text{Ker}(\chi) = 1$.

Assume that χ^2 is a multiple of an irreducible. In [2] it is proved that if $\chi, \psi \in \text{Irr}(G)$ are faithful characters of a finite p -group G , and the product $\chi\psi$ is a multiple of an irreducible, then χ and ψ vanish outside the center of G . Since $\chi^{(2)}$ is a constituent of χ^2 and χ^2 is a multiple of an irreducible, (i) follows.

We may assume now that χ^2 is not a multiple of an irreducible. In [1] it is proved that if $\chi, \psi \in \text{Irr}(G)$ are faithful characters of a finite p -group, where p is odd, either $\chi\psi$ is a multiple of an irreducible or $\chi\psi$ has at least $\frac{p+1}{2}$ distinct irreducible constituents. By Proposition 2.5 we have that χ^2 does not have any linear constituent. Since $\chi^2(1) = p^2$ and χ^2 is not a multiple of an irreducible, it follows that all the constituents of χ^2 have degree p . By Theorem A we have that χ^2 has a unique irreducible constituent that appears with odd multiplicity. Since χ^2 has at least $\frac{p+1}{2}$ distinct irreducible constituents, $\chi^{(2)}$ is the unique irreducible constituent that appears with odd multiplicity and

$$\chi(1)^2 = p^2 = p + 2\left(\frac{p-1}{2}\right)p,$$

it follows that χ^2 has to have exactly $\frac{p+1}{2}$ distinct irreducible constituents and so (ii) follows. □

Proof of Theorem C. Since any nilpotent subgroup is isomorphic to the direct sum of its p -Sylow subgroups, Theorem C follows from Lemma 2.9. □

Remark 2.10. Let p and q odd prime numbers such that $q = rp + 1$, for some integer r , with $1 < r < \frac{p+1}{2}$. Let G be a nonabelian group of order pq . We can check that G has exactly p distinct linear characters and r distinct irreducible characters of degree p .

Let $\chi \in \text{Irr}(G)$ with $\chi(1) = p$. Since G has odd order, then χ^2 does not have any linear constituent. Thus χ^2 is the sum of at most r distinct irreducible characters of G , where $r < \frac{p+1}{2}$. We can check that χ^2 is not a multiple of an irreducible. Therefore Theorem C does not remain true with the weaker hypothesis that G is supersolvable of odd order.

REFERENCES

- [1] E. Adan-Bante, Products of characters and finite p -groups, J. Algebra 277, 236-255.
- [2] E. Adan-Bante, M. Loukaki, A. Moreto, Homogeneous products of characters, J. Algebra 274, 587-593.
- [3] I. M. Isaacs, Character Theory of Finite Groups. New York-San Francisco-London: Academic Press 1976.

UNIVERSITY OF SOUTHERN MISSISSIPPI GULF COAST, 730 EAST BEACH BOULEVARD, LONG BEACH MS 39560

E-mail address: Edith.Bante@usm.edu